



**QUEEN'S
UNIVERSITY
BELFAST**

Tolerance to sedative drugs in PICU: Can it be moderated or is it immutable?

Wolf, A. R., Blackwood, B., & Anderson, B. (2016). Tolerance to sedative drugs in PICU: Can it be moderated or is it immutable? *Intensive Care Medicine*, 42 (2), 278–281. <https://doi.org/10.1007/s00134-015-4119-0>

Published in:
Intensive Care Medicine

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
© [2015] Springer
The final publication is available at Springer via <http://link.springer.com/article/10.1007%2Fs00134-015-4119-0>

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Intensive Care Medicine

Tolerance to sedative drugs in PICU: Can it be moderated or is it immutable?

--Manuscript Draft--

Manuscript Number:	
Full Title:	Tolerance to sedative drugs in PICU: Can it be moderated or is it immutable?
Article Type:	Editorial
Funding Information:	
Corresponding Author:	Andy Wolf, M.B., BChir(Cantab), M. D(Bristol), FRLA our lady's children hospital dublin, IRELAND
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	our lady's children hospital
Corresponding Author's Secondary Institution:	
First Author:	Andy Wolf, M.B., BChir(Cantab), M. D(Bristol), FRLA
First Author Secondary Information:	
Order of Authors:	Andy Wolf, M.B., BChir(Cantab), M. D(Bristol), FRLA
	Bronagh blackwood
	brian anderson
Order of Authors Secondary Information:	
Author Comments:	
Suggested Reviewers:	

[Click here to view linked References](#)

Tolerance to sedative drugs in PICU: Can it be moderated or is it immutable?

Andrew Wolf
Department of Anaesthesia and Intensive Care
Our Lady's Children's Hospital, Crumlin,
Dublin 12
Ireland

Bronagh Blackwood
School of Medicine, Dentistry and Biomedical Sciences
Queens University, University Road,
Belfast
Northern Ireland BT7 1NN

Brian Anderson
Departments of Anaesthesiology
University of Auckland FHMS
Park Road Grafton,
Auckland 1023
New Zealand

Address for Correspondence:
Professor Andrew R Wolf
awolfbch@aol.com, andrew.wolf@nhs.net

1 Nienke Vet and colleagues have evaluated the influence of daily sedation interruptions
2 (DSI) in PICU. This was ambitious: bias is hard to avoid and recruitment rates in such
3 studies are poor, which ultimately restrict the strength of conclusions. Nevertheless,
4 their results suggest that if behavioral tools already drive sedation delivery, then DSI
5 will not substantially influence the duration of ventilation. These findings are consistent
6 with recent reports from adult intensive care (1).
7
8
9
10
11
12
13
14
15

16 A pause in drug delivery allows plasma concentration to fall followed by reduction in
17 effect site concentration and subsequent arousal. However, the lag time can be
18 significant (Figure 1a). Recovery may be further delayed by the presence and influence
19 of midazolam's alpha-hydroxy metabolite that has half the activity of the parent drug
20 (2). Other drugs used for sedation also have influence; a three-drug combination (e.g.,
21 midazolam, propofol, and alfentanil) can triple the duration of effect compared with
22 propofol alone (3). In the current study additional sedative and analgesic drugs were
23 used "as required". This may have contributed to the lack of impact from midazolam
24 interruption. Variation in recovery from sedation is complicated by the high prevalence
25 of renal failure, hepatic failure, and concomitant administration of CYP3A inhibitors in
26 PICU patients. Sedation requirements vary widely not only with age, diagnosis and
27 clinical state, but also between similar patients.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 Sedation has parallels with innovations in postoperative pain control in children:
46 intermittent drug use was replaced by continuous infusions, then patient and nurse
47 controlled analgesia (PCA/NCA) with pain score monitoring. Further optimization
48 combined low dose continuous infusions modulated by PCA/NCA. PICU sedation is in
49 need of similar improvement but can this impact on patient outcomes? Practitioners of
50 anaesthesia are also aware of emergence delirium complicating recovery. Could a
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 similar phenomenon complicate midazolam interruption, necessitating reintroduction
2 of the drug prematurely?
3
4

5
6 The value of DSI (4,5,6) remains unclear. The results of this current study could simply
7 reflect the strong history in the Netherlands of good sedation practice including a
8 relative sparing use of sedatives. This is supported by the relatively low dose of
9 midazolam used compared to a previous paediatric study (6). One conclusion is that
10 recovery from sedation relates to the overall cumulative “sedation burden” and that
11 minimizing the exposure with effective behavioral scoring linked to delivery, optimizes
12 recovery irrespective of DSI and other factors.
13
14
15
16
17
18
19
20
21
22
23
24

25 Interventions such as those described by Vet and colleagues are complex. They consist
26 of multiple linked elements: (7) context and setting (workload, resources, staffing); ICU
27 staff characteristics (skill mix, training,) and clinical processes (complexity of protocols,
28 algorithms, decision-making and perceived risk) (8). Understanding these components
29 is critical to interpreting and generalizing study findings (9). This study does not
30 provide contextual information that may help explain the results. An indication of the
31 degree of compliance and nursing perspectives about the DSI may assist with
32 interpreting the findings. Qualitative studies exploring non-adherence to DSI have
33 highlighted a lack of nursing acceptance due to patient agitation and the subsequent risk
34 of adverse events associated with more wakeful patients (11, 12). The UK Medical
35 Research Council recommends that a process evaluation should accompany trials of
36 complex interventions. (9).
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 Can tolerance be manipulated by modifying practice? Techniques that should be
55 considered include: avoiding drugs particularly associated with tolerance such as
56 midazolam, moderating early exposure to high doses of analgesics and sedatives to
57
58
59
60
61
62
63
64
65

1 prevent acute tolerance, drug-sparing strategies such as drug cycling or polypharmacy
2 mixtures, and returning to low dose propofol, a drug that has been legislated out of PICU
3 in many countries. Midazolam remains the most popular sedative despite association
4 with withdrawal phenomena of up to 35% (13). The frequency and severity of tolerance
5 and withdrawal is related to the cumulative “drug burden” and higher doses (≥ 300
6 mcg/kg/h) (14). A sedated patient is thought to be easier to nurse (15) and even in this
7 current study children were sedated with mean midazolam doses (183-240 mcg/kg/h)
8 to the deeper end of the sedation scale. There are alternatives to midazolam. Alpha-2
9 agonists can provide effective sedation either as a straight midazolam replacement or as
10 a supporting drug (16,17). Currently there are no substantial data to determine if other
11 drug combinations can reduce tolerance and accelerate recovery.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 Acute exposure to high doses of high-efficacy short-acting opioids (e.g. fentanyl,
28 remifentanyl) is linked to accelerated drug tolerance (18) and delay in recovery. Fast
29 track paediatric cardiac surgery has moved away from high doses fentanyl 100 mcg/kg
30 or more to 10-15 mcg/kg over the perioperative period resulting in earlier extubation
31 and accelerated recovery. There are potentially transferable lessons from this
32 experience that could reduce tolerance in PICU patients. Drug delivery must reflect the
33 age and context dependent pharmacokinetics. For example while drug delivery in the
34 young infant needs to be high during the loading phase, downward adjustment is
35 necessary during maintenance, reflecting reduced elimination compared to the older
36 infant (Figure 1b). Unfortunately, behavioral scoring may be impossible in this early
37 phase if neuromuscular blocking drugs are used, and higher doses than necessary may
38 be continued. Rotating drug sedation and analgesic regimens, or using non-
39 pharmacological strategies to maintain comfort have been used to try to limit drug
40 requirements. While this approach makes sense there are no data to support it.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 The SLEEPS trial (16) demonstrated that neither morphine/midazolam nor
2 morphine/clonidine alone could always provide complete sedation. A third agent was
3
4 often required. The choice of the third agent may be important: in the current study
5
6 multiple rescue drugs were used. While propofol as a major sedative agent has been
7
8 eliminated from use in PICU due to fears of propofol infusion syndrome, it continues to
9
10 be used cautiously by some, even in countries where the drug is officially discouraged.
11
12 Low-dose infusion (0-4 mg/kg/h) with careful surveillance for accumulation and lactic
13
14 acidosis deserves to be reconsidered as a third-line drug.
15
16

17
18
19
20 Sedation is often treated as a necessary evil in PICU: the primary disease and its
21
22 treatment is naturally the main focus while sedation is managed generically. During
23
24 recovery the secondary problems associated with sedation (e.g., nosocomial infection,
25
26 poor gut motility and behavioral change) become lost within the disease and general
27
28 PICU experience. Under-sedation and over-sedation are both harmful: minimizing
29
30 sedation exposure and their adverse effects are important. Optimized matching of
31
32 delivery to sedation requirement provides another marginal gain in the critically ill child
33
34 that can contribute to improving patient outcomes.
35
36
37

38
39 On behalf of all authors, the corresponding author states that there is no conflict of
40
41 interest.
42
43
44
45

46 **References**

- 47
48
49
50
51 1. Burry L, Rose L, McCullagh IJ, *et al.* Daily sedation interruption versus no daily
52 sedation interruption for critically ill adult patients requiring invasive
53 mechanical ventilation. *Cochrane Database Syst Rev* 2014; 7: CD009176.
54
55
56 2. Mandema JW, Tuk B, van Steveninck AL, *et al.* Pharmacokinetic-
57 pharmacodynamic modeling of the central nervous system effects of midazolam
58 and its main metabolite alpha-hydroxymidazolam in healthy volunteers. *Clin*
59 *Pharm Ther* 1992; 51: 715-728.
60
61
62
63
64
65

- 1 3. Minto CF, Schnider TW, Short TG, *et al.* Response surface model for anesthetic
2 drug interactions. *Anesthesiology* 2000; **92**: 1603-1616.
- 3
- 4 4. Mehta s, Burry L, Cook D Daily sedation interruption in mechanically
5 ventilated critically ill patients cared for with a sedation protocol: a randomized
6 controlled trial. *JAMA* 2012; 308: 1985 - 1992
- 7
- 8 5. Verlaad CW, Heesen GP, Vet NJ et al. Randomised controlled trial of daily
9 interruption of sedatives in critically ill children. *Pediatric Anesthesia* 2014 24;
10 151-156
- 11
- 12 6. Gupta K, Gupta VK, Jayashree M et al Randomised controlled trial of
13 interrupted versus sedative infusions in ventilated children. *Pediatr Crit Care*
14 2012 Med 13: 131-135.
- 15
- 16 7. Blackwood B. Methodological issues in evaluating complex health care
17 interventions. *Journal of Advanced Nursing* 2006, 54 (5): 612-622
- 18
- 19 8. Rose L, Blackwood B, Burns SM, Frazier SK, Egerod I. International
20 perspectives on the influence of structure and process of weaning from
21 mechanical ventilation. *American Journal of Critical Care*, 2011; 20 (1), e10-e18
22 doi: 10.4037/ajcc2011xxx
- 23
- 24 9. Moore GF, Audrey S, Barker M, Bond L, Bonell C, Hardeman W, Moore L,
25 O’Cathain A, Tinati T, Wight D, Baird J. Process evaluation of complex
26 interventions: Medical Research Council guidance. *BMJ* 2015;350:h1258
27 doi:10.1136/bmj.h1258
- 28
- 29 10. Shehabi Y & Weisbrodt L. Daily sedation interruption; a glass half empty.
30 *Anaesthesia and Intensive Care* 2011, 39, 339–341.
- 31
- 32 11. Everingham K, Fawcett T, Walsh T. Targeting’ sedation: the lived experience
33 of the intensive care nurse. *Journal of Clinical Nursing*, 2014, 23, 694–703
- 34
- 35 12. Tanios M, de Wit M, Epstein S & Devlin . Perceived barriers to the use of
36 sedation protocols and daily sedation interruption: a multidisciplinary survey.
37 *Journal of Critical Care* 2009, 24, 66–73.
- 38
- 39 13. Hughes J, Gill A, Leach HJ et al. A prospective study of the adverse effects of
40 midazolam on withdrawal in critically ill children. *Acta Paediatr*
41 1994;83(11):1194-9.
- 42
- 43 14. Jenkins IA, Playfor SD, Bevan C, et al. Current United Kingdom sedation
44 practice in pediatric intensive care. *Paediatric Anaesthesia* 2007; 17 (7): 675-83.
- 45
- 46 15. Sneyers B, Laterre PF, Bricq E, *et al.* What stops us from following sedation
47 recommendations in intensive care units? A multicentric qualitative study. *J Crit*
48 *Care* 2014; **29**: 291-297.
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

- 1 16. Wolf A, McKay A, Spowart C et al Prospective multi-centre randomised,
2 double-blind, equivalence study comparing clonidine and midazolam as
3 intravenous sedative agents in critically ill children. The SLEEPS Study (Safety
4 ProfiLe, Efficacy and Equivalence in Paediatric intensive care Sedation)
5 Health Technology Assessment 2014. 18(71):1-212, 2014 Dec
6
7
8 17. Achuff BJ; Nicolson SC; Elci OU; Zuppa AF. Intraoperative dexmedetomidine
9 reduces postoperative mechanical ventilation in infants after open heart surgery.
10 Pediatric Critical Care Medicine 2015. 16(5):440-7.
11
12
13 18. Crawford MW; Hickey C; Zaarour C; Howard A; Naser B. Development of
14 Acute opioid tolerance during remifentanyl infusion for pediatric scoliosis
15 surgery. Anesthesia and Analgesia 2006 102(6):1662-7.
16
17
18
19
20
21
22
23
24
25

26 **Figure 1a.** Plasma concentrations and effect in a 2-year-old child given protocol midazolam
27
28 bolus (0.1 mg/kg) and infusion changes (100 mcg/kg/h step changes) every 30 min to achieve
29
30 sedation. Infusion was stopped at 180 min. Sedation recovery lags behind the decline in
31
32 plasma concentration. Amplitudes in the 11.5-30 Hz (beta) frequency band were used as an
33
34 EEG effect measure. Pharmacodynamic parameter estimates were from Mandema J et al.
35
36 Pharmacokinetic-pharmacodynamic modeling of the central nervous system effects of
37
38 midazolam and its main metabolite alpha-hydroxymidazolam in healthy volunteer. Clin
39
40 Pharm Ther 1992;51:715-28
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65






Figure 1b. Plasma concentrations and effect in a neonate given protocol midazolam bolus (0.1 mg/kg) on two early occasions (5 min interval) to achieve sedation. Plasma concentration declines slowly because of slow clearance. Sedation recovery lags way behind the decline in plasma concentration even though a maintenance infusion was not even given. Pharmacodynamic parameter estimates were from Mandema J et al. Pharmacokinetic-pharmacodynamic modeling of the central nervous system effects of midazolam and its main metabolite alpha-hydroxymidazolam in healthy volunteer. Clin Pharm Ther 1992;51:715-28



[Click here to view linked References](#)



[Click here to view linked References](#)

